

TECHNICAL DATA SHEET

APC Anti-Human CD4 (SK3)

Catalog Number: 20-0047

PRODUCT INFORMATION

Contents: APC Anti-Human CD4 (SK3)

Isotype: Mouse IgG1, kappa

Concentration: 5 μ L (0.06 μ g)/test


Clone: SK3

Reactivity: Human

Use By: 12 months from date of receipt

Storage Conditions: 2-8°C protected from light

Formulation: 10 mM NaH₂PO₄, 150 mM NaCl, 0.09% NaN₃, 0.1% gelatin, pH7.2



Human peripheral blood lymphocytes were stained with 5 μ L (0.06 μ g) APC Anti-Human CD4 (20-0047) (solid line) or 0.06 μ g APC Mouse IgG1 isotype control (dashed line).

DESCRIPTION

The SK3 antibody reacts with human CD4, a 59 kDa protein which acts as a co-receptor for the T cell receptor (TCR) in its interaction with MHC Class II molecules on antigen-presenting cells. The extracellular domain of CD4 binds to the beta-2 domain of MHC Class II, while its cytoplasmic tail provides a binding site for the tyrosine kinase lck, facilitating the signaling cascade that initiates T cell activation. CD4, and co-receptors CCR5 and CXCR4, may also be utilized by HIV-1 to enter T cells. Human CD4 is typically expressed on thymocytes, some mature T cell populations such as Th17 and T regulatory (Treg) cells, as well as on dendritic cells. The SK3 antibody is widely used as a phenotypic marker for human CD4 expression, and has been reported to be cross-reactive with Rhesus and Cynomolgus CD4. This antibody does not block binding of alternative clone RPA-T4, suggesting that they recognize different epitopes.

PREPARATION & STORAGE

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

APPLICATION NOTES

This antibody preparation has been pre-titrated and quality-tested for flow cytometry using an appropriate cell type. The antibody has been diluted for use at 5 μ L per test, defined as the amount of antibody that will stain a cell sample in a final volume of approximately 100 μ L. The number of cells within a sample should be determined empirically, but typically ranges between 1x10⁵ to 1x10⁸ cells.

REFERENCES

- Evans RL, Wall DW, Platsoucas CD, Siegal FP, Fikrig SM, Testa CM and Good RA. 1981. Proc Natl Acad Sci U S A. 78(1): 544-548.
- Sattentau QJ, Dalgleish AG, Weiss RA and Beverley PC. 1986. Science. 234(4780): 1120-1123.
- Bernard A, Boumsell L and Hill C. In: Bernard A, Boumsell L, Dausset J, Milstein C, Schlossman SF, ed. Leucocyte Typing. New York, NY: Springer-Verlag; 1984: 9-108. (Flow Cytometry)
- Heninger AK, Theil A, Wilhelm C, Petzold C, Huebel N, Kretschmer K, Bonifacio E and Monti P. 2012. J Immunol. 189(12): 5649-5648. (Flow Cytometry)
- Yoshino N, Ami Y, Terao K, Tashiro F and Honda M. 2000. Exp. Anim. 49(2): 97-110. (Flow Cytometry - Cynomolgus)
- Lafont BAP, Gloeckler L, D'Hautcourt JL, Gut JP and Aubertin AM. 2000. Cytometry 41: 193-202. (Flow Cytometry - Rhesus)

Tonbo Biosciences tests all antibodies by flow cytometry. Citations are provided as a resource for additional applications that have not been validated by Tonbo Biosciences. Please choose the appropriate format for each application and consult Materials and Methods sections for additional details about the use of any product in these publications.

For Research Use Only.

Not for use in diagnostic or therapeutic procedures. Not for resale. Not for distribution without written consent. Tonbo Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Tonbo Biosciences, Tonbo Biosciences Logo and all other trademarks are the property of Tonbo Biotechnologies Corporation. © 2013 Tonbo Biosciences.