

TECHNICAL DATA SHEET

# APC Anti-Mouse CD252 (OX40 Ligand) (RM134L)

Catalog Number: 20-5905

## PRODUCT INFORMATION

**Contents:** APC Anti-Mouse CD252 (OX40 Ligand) (RM134L)

**Isotype:** Rat IgG2b, kappa

**Concentration:** 0.2 mg/mL


**Clone:** RM134L

**Reactivity:** Mouse

**Use By:** 12 months from date of receipt

**Storage Conditions:** 2-8°C

**Formulation:** 10 mM NaH<sub>2</sub>PO<sub>4</sub>, 150 mM NaCl, 0.09% Na<sub>3</sub>N, 0.1% gelatin, pH7.2



C57Bl/6 splenocytes were stimulated with anti-IgM and anti-CD40 for 4 days. Cells were then stained with 0.25 ug APC Anti-Mouse CD252 (20-5905) (solid line) or 0.25 ug APC Rat IgG2b isotype control (dashed line).

## DESCRIPTION

The RM134L antibody recognizes CD252, also known as OX-40 Ligand or CD134 Ligand, a member of the TNF superfamily that is present on the surface of antigen presenting cells and activated B lymphocytes. The OX-40 Ligand interacts with OX-40 (CD134) which is expressed primarily on activated T cells. This costimulatory interaction leads to increased proliferation and IL-2 production responses of activated T cells, and at the same time enhances proliferation and immunoglobulin secretion by activated B cells. The RM134L antibody is useful for flow cytometric detection of CD252 on stimulated mouse splenocytes. It has also been reported to block the costimulatory activity of OX-40 Ligand. Please choose the appropriate format for each application.

## PREPARATION & STORAGE

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

## APPLICATION NOTES

This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). The amount of antibody required for optimal staining of a cell sample should be determined empirically in your system.

## REFERENCES

- Akiba H, Oshima H, Takeda K, Atsuta M, Nakano H, Nakajima A, Nohara C, Yagita H and Okumura K. 1999 J Immunol. 162(12): 7058-7066. (Flow cytometry, in vitro blocking)
- Stuber E, Neurath M, Calderhead D, Fell HP and Strober W. 1995 Immunity. 2(5): 507-521.
- Sibilano R, Frossi B, Suzuki R, D'Inca F, Gri G, Piconese S, Colombo MP, Rivera J and Pucillo CE. 2012 Journal of Allergy Clin Immunol. 130(3): 751-760. (Blocking)
- van der Merwe M, Abdelsamed HA, Seth A, Ong T, Vogel P and Pillai AB. 2013 J Immunol. 191(11): 5764-5776 (Flow cytometry)

Tonbo Biosciences tests all antibodies by flow cytometry. Citations are provided as a resource for additional applications that have not been validated by Tonbo Biosciences. Please choose the appropriate format for each application and consult Materials and Methods sections for additional details about the use of any product in these publications.

For Research Use Only.

Not for use in diagnostic or therapeutic procedures. Not for resale. Not for distribution without written consent. Tonbo Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Tonbo Biosciences, Tonbo Biosciences Logo and all other trademarks are the property of Tonbo Biotechnologies Corporation. © 2013 Tonbo Biosciences.