

TECHNICAL DATA SHEET

# Biotin Anti-Mouse IFN gamma (XMG1.2)

Catalog Number: 30-7311

## PRODUCT INFORMATION

**Contents:** Biotin Anti-Mouse IFN gamma (XMG1.2)

**Isotype:** Rat IgG1, kappa

**Concentration:** 0.5 mg/mL


**Clone:** XMG1.2

**Reactivity:** Mouse

**Use By:** 12 months from date of receipt

**Storage Conditions:** 2-8°C

**Formulation:** 10 mM NaH<sub>2</sub>PO<sub>4</sub>, 150 mM NaCl, 0.09% Na<sub>3</sub>N, pH7.2



C57Bl/6 splenocytes were stimulated with PMA and Ionomycin and then stained with APC Anti-Mouse CD3e (20-0031), followed by intracellular staining with 0.25 ug Biotin Anti-Mouse IFN gamma (30-7311) (right panel) or 0.25 ug Biotin Rat IgG1 isotype control (left panel), detected with Streptavidin PE.

## DESCRIPTION

The XMG1.2 antibody is specific for mouse Interferon-gamma (IFN-g), a 20 kDa type II cytokine known for its central roles in protection against bacterial or viral pathogens and for its anti-tumor properties. IFN-g is secreted by several types of immune cells which allow the cytokine to modulate innate immunity when secreted by NK and NKT cells, and to function in support of adaptive immunity when secreted by Th1 and CD8+ T cells (CTLs). The XMG1.2 antibody is suitable for detection of intracellular IFN-g protein by flow cytometry. This format can be used for quantitative analysis of the secreted protein by ELISA when paired with an appropriate capture antibody. This clone has been reported for neutralization of the functional activity of IFN-g in a variety of assays (use format suitable for functional assays).

## PREPARATION & STORAGE

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

## APPLICATION NOTES

This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). The amount of antibody required for optimal staining of a cell sample should be determined empirically in your system.

## REFERENCES

Choudhry N, Petry F, van Rooijen N, and McDonald V. 2012. *J. of Infect. Disease.* 206: 117-124. (in vivo neutralization) Cobb D and Smeltz RB. 2012. *J. Immunol.* 188: 3766-3773. (in vitro neutralization) Brown DM, Lee S, Garcia-Hernandez M, and Swain SL. 2012. *J. Virol.* 86: 6792-6803. (ELISpot - detection) Yu H, Karunakaran KP, Jiang X, Shen C, Andersen P, and Brunham RC. 2012. *Infect. Immun.* 80: 1510-1518. (ELISpot -detection) Kwon M-J, Ma J, Ding Y, Wang R, and Sun Z. 2012. *J. Immunol.* 188: 5887-5897. (in vitro induction of Th2 polarization) Barr TA, Shen P, Brown S, Lampropoulou V, Roch T, Lawrie S, Fan B, O'Connor RA, Anderton SM, Bar-Or Am Fillatreau S, and Gray D. 2012. *J. Exp. Med.* 209: 1001-1010. (Flow cytometry) Cardona AE, Restrepo BI, Jaramillo JM, and Teale JM. 1999. *J. Immunol.* 162: 995-1002. (Immunohistochemistry – frozen tissue) Kupfer A, Mosmann TR, and Kupfer H. 1991. *Proc. Natl. Acad. Sci.* 88: 775-779. (Immunofluorescence microscopy)

Tonbo Biosciences tests all antibodies by flow cytometry. Citations are provided as a resource for additional applications that have not been validated by Tonbo Biosciences. Please choose the appropriate format for each application and consult Materials and Methods sections for additional details about the use of any product in these publications.

For Research Use Only.

Not for use in diagnostic or therapeutic procedures. Not for resale. Not for distribution without written consent. Tonbo Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Tonbo Biosciences, Tonbo Biosciences Logo and all other trademarks are the property of Tonbo Biotechnologies Corporation. © 2013 Tonbo Biosciences.