

TECHNICAL DATA SHEET

PE Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2)

Catalog Number: 50-5321

PRODUCT INFORMATION

Contents: PE Anti-Mouse MHC Class II (I-A/I-E) (M5/114.15.2)


Isotype: Rat IgG2b, kappa

Concentration: 0.2 mg/mL

Clone: M5/114.15.2

Reactivity: Mouse

Formulation: 10 mM NaH₂PO₄, 150 mM NaCl, 0.09% Na₃N, 0.1% gelatin, pH7.2



C57Bl/6 splenocytes were stained with 0.06 ug Anti-Mouse MHC Class II PE (50-5321) (solid line) or 0.06 ug Rat IgG2b PE isotype control (dashed line).

DESCRIPTION

The M5/114.15.2 antibody reacts with mouse MHC Class II alloantigens I-Ab, I-Ad, I-Aq, I-Ed, and I-Ek, as well as being cross-reactive with mouse cells of H-2p and H-2r haplotype. MHC Class II is widely expressed by mouse immune cells bearing these alloantigens, including T and B cells, monocytes, macrophages, and dendritic cells. The antibody does not react with the following alloantigens: I-Af, I-Ak, I-As, or NOD H-2g. The M5/114.15.2 antibody may be used for analysis of mouse cells expressing MHC Class II alloantigens as described. Please note that the M5/114.15.2 clone may also be referred to as M5/114 in the literature.

PREPARATION & STORAGE

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

APPLICATION NOTES

This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). The amount of antibody required for optimal staining of a cell sample should be determined empirically in your system.

REFERENCES

Staeheil F, Ludigs K, Heinz LX, Segin-Estevez Q, Ferrero I, Braun M, Schroder K, Rebsamen M, Tardivel A, Mattmann C, MacDonald HR, Romero P, Reith W, Guarda G, and Tschopp J. 2012. *J. Immunol.* 188: 3820-3828. (Flow cytometry). Parra D, Rieger AM, Li J, Zhang Y-A, Randall LM, Hunter CA, Barreda DR, and Sunyer JO. 2012. *J. Leukoc. Biol.* 91:525-536. (in vitro blocking, Flow cytometry). Scarlett UK, Rutkowski MR, Rauwerdink AM, Fields J, Escovar-Fadul X, Baird J, Cubillos-Ruiz JR, Jacobs AC, Gonzalez JL, Weaver J, Fiering S, and Conejo-Garcia JR. 2012. *J. Exp. Med.* 209: 495-506. (Immunofluorescence microscopy – frozen tissue). Chen M, Felix K, and Wang J. 2011. *J. Immunol.* 187: 5684-5692. (in vitro blocking). Busman-Sahay K, Sargent E, Harton JA, and Drake JR. 2011. *J. Immunol.* 186:6710-6717. (Immunoprecipitation). Ohmura-Hoshino M, Matsuki Y, Aoki M, Goto E, Mito M, Uematsu M, Hakiuchi T, Hotta H, and Ishido S. 2006. *J. Immunol.* 177:341-354. (Immunofluorescence microscopy – frozen tissue, Immunoprecipitation). Li C, Siemasko K, Clark MR, and Song W. 2002. *Int. Immunol.* 14: 1179-1191. (Western Blot, Immunoelectron microscopy).