

TECHNICAL DATA SHEET

PE-Cyanine7 Anti-Mouse CD19 (1D3)

Catalog Number: 60-0193

PRODUCT INFORMATION

Contents: PE-Cyanine7 Anti-Mouse CD19 (1D3)

Isotype: Rat IgG2a, kappa

Concentration: 0.2 mg/mL


Clone: 1D3

Reactivity: Mouse

Use By: 6 months from date of receipt

Storage Conditions: 2-8°C protected from light

Formulation: 10 mM NaH₂PO₄, 150 mM NaCl, 0.09% NaN₃, 0.1% gelatin, pH7.2



C57Bl/6 splenocytes were stained with 0.25 ug PE-Cy7 Anti-Mouse CD19 (60-0193) (solid line) or 0.25 ug PE-Cy7 Rat IgG2a isotype control (dashed line).

DESCRIPTION

The 1D3 antibody reacts with mouse CD19, a 95 kDa glycoprotein which acts as a co-receptor, along with CD21 and CD81, in support of the functional B cell receptor (BCR). This complex provides antigen-specific recognition and subsequent activation of B cells to proliferate and differentiate into antibody-secreting cells (plasma cells) or memory B cells, which are crucial for secondary antigen encounter. CD19 is a lineage-differentiation marker, as its expression is detectable at the earliest B cell stages, through development, and is finally lost upon transition to mature plasma cells. The 1D3 antibody is widely used as a phenotypic marker for CD19 expression on B cells, as well as on dendritic cell subsets.

PREPARATION & STORAGE

This monoclonal antibody preparation was purified from tissue culture supernatant via affinity chromatography. For In Vivo Ready™ (IVR) products, each preparation is also evaluated for endotoxin levels using the LAL assay. It is recommended to store the product undiluted at 4°C. Do not freeze.

APPLICATION NOTES

This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). Please refer to the figure legend for the optimal concentration used to stain the tissue shown. We recommend titrating the antibody under your specific conditions to determine the optimal concentration of antibody needed in your experimental system.

REFERENCES

Ghoshn EEB, Yamamoto R, Hamanada S, Yang Y, Herzenberg LA, Nakauchi H, and Herzenberg LA. 2012. Proc. Natl. Acad. Sci. 109:5394-5398. (Flow cytometry)
 Raghavan S, Ostberg AK, Flach C-F, Ekman A, Blomquist M, Czerkinsky C, and Holmgren J. 2010. Infect. Immun. 78(10):4251-4260. (Immunohistochemistry – acetone fixed tissue)
 Togayachi A, Kozono Y, Ikehara Y, Ito H, et al. 2010. Proc. Natl. Acad. Sci. 107:11900-11905. (Immunoprecipitation)
 Poitrasson-Riviere M, Bienvenu B, Le Campion A, Becourt C, Martin B, and Lucas B. 2008. J. Immunol. 180:7294-7304. (Immunohistochemistry – frozen tissue)
 Lee Y, Haas KM, Gor DO, Ding X, Karp DR, Greenspan NS, Poe JC, and Tedder TF. 2005. J. Immunol. 175:8011-8023. (Immunoprecipitation)
 Bobbitt KR and Justement LB. 2000. J. Immunol. 165: 5588-5596. (in vitro stimulation, Immunoprecipitation)

Tonbo Biosciences tests all antibodies by flow cytometry. Citations are provided as a resource for additional applications that have not been validated by Tonbo Biosciences. Please choose the appropriate format for each application and consult Materials and Methods sections for additional details about the use of any product in these publications.

For Research Use Only.

Not for use in diagnostic or therapeutic procedures. Not for resale. Not for distribution without written consent. Tonbo Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Tonbo Biosciences, Tonbo Biosciences Logo and all other trademarks are the property of Tonbo Biotechnologies Corporation. © 2013 Tonbo Biosciences.