

TECHNICAL DATA SHEET

# PE-Cyanine7 Anti-Human HLA-DR (LN3)

Catalog Number: 60-9956

## PRODUCT INFORMATION

**Contents:** PE-Cyanine7 Anti-Human HLA-DR (LN3)

**Isotype:** Mouse IgG2b, kappa

**Concentration:** 5ul (0.125ug)/test


**Clone:** LN3

**Reactivity:** Human

**Use By:** 6 months from date of receipt

**Storage Conditions:** 2-8°C protected from light

**Formulation:** 10 mM NaH<sub>2</sub>PO<sub>4</sub>, 150 mM NaCl, 0.09% NaN<sub>3</sub>, 0.1% gelatin, pH7.2



Human peripheral blood monocytes were stained with 5 uL (0.125 ug) PE-Cyanine7 Anti-Human HLA-DR (60-9956) (solid line) or 0.125 ug PE-Cyanine7 Mouse IgG2b isotype control (dashed line).

## DESCRIPTION

The LN3 antibody reacts with a member of the human MHC Class II antigens, HLA-DR. The HLA-DR antigen is expressed on human antigen presenting cells including B lymphocytes, monocytes, macrophages, dendritic cells, and activated T lymphocytes. HLA-DR is a dimeric protein composed of alpha and beta subunits and is involved in the presentation of peptide antigens to CD4+ T cells.

## PREPARATION & STORAGE

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

## APPLICATION NOTES

This antibody preparation has been pre-titrated and quality-tested for flow cytometry using an appropriate cell type. The antibody has been diluted for use at 5 µL per test, defined as the amount of antibody that will stain a cell sample in a final volume of approximately 100 µL. The number of cells within a sample should be determined empirically, but typically ranges between 1x10<sup>5</sup> to 1x10<sup>8</sup> cells.

## REFERENCES

Eckert F, Schmid U. 1989. Arch Dermatol. 125(11):1518-1524. Swanson PE, Wick MR. 1990. Mod Pathol. 3(2):113-119. Shin SS, Sheibani K, Kezirian J, Nademanee A, Forman SJ, Lee SK, Winberg CD. 1992. Hum Pathol. 23(6):686-694. Sugita S, Iwasaki Y, Makabe K, Kimura T, Futagami T, Suegami S, Takahashi M. 2016. Stem Cell Reports. 11;7(4):619-634.

Tonbo Biosciences tests all antibodies by flow cytometry. Citations are provided as a resource for additional applications that have not been validated by Tonbo Biosciences. Please choose the appropriate format for each application and consult Materials and Methods sections for additional details about the use of any product in these publications.

For Research Use Only.

Not for use in diagnostic or therapeutic procedures. Not for resale. Not for distribution without written consent. Tonbo Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Tonbo Biosciences, Tonbo Biosciences Logo and all other trademarks are the property of Tonbo Biotechnologies Corporation. © 2013 Tonbo Biosciences.