

TECHNICAL DATA SHEET

# PerCP-Cyanine5.5 Anti-Mouse CD134 (OX-86)

Catalog Number: 65-1341

## PRODUCT INFORMATION

**Contents:** PerCP-Cyanine5.5 Anti-Mouse CD134 (OX-86)

**Isotype:** Rat IgG1, kappa

**Concentration:** 0.2 mg/mL


**Clone:** OX-86

**Reactivity:** Mouse

**Use By:** 6 months from date of receipt

**Storage Conditions:** 2-8°C protected from light

**Formulation:** 10 mM NaH<sub>2</sub>PO<sub>4</sub>, 150 mM NaCl, 0.09% NaN<sub>3</sub>, 0.1% gelatin, pH7.2



C57Bl/6 splenocytes were stimulated for 3 days with ConA and then stained with APC Anti-Mouse CD3 (20-0031) and 0.25 ug PerCP-Cyanine5.5 Anti-Mouse CD134 (65-1341) (right panel) or 0.125 ug PerCP-Cyanine5.5 Rat IgG1 (left panel).

## DESCRIPTION

The OX-86 monoclonal antibody reacts with mouse CD134, a 50kD type 1 integral membrane glycoprotein and a member of the TNF receptor superfamily. CD134 is also known as OX40 and is expressed on activated CD4 and CD8 T cells. CD134 is the receptor for CD252 (OX40 ligand) and the interaction of this receptor-ligand pair is implicated in T-dependent humoral responses as well as numerous aspects of T cell expansion, survival and memory. Additionally, CD134 seems to play a significant role in the pathogenesis of some autoimmune diseases.

## PREPARATION & STORAGE

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

## APPLICATION NOTES

This antibody preparation has been quality-tested for flow cytometry using mouse spleen cells, or an appropriate cell type (where indicated). Please refer to the figure legend for the optimal concentration used to stain the tissue shown. We recommend titrating the antibody under your specific conditions to determine the optimal concentration of antibody needed in your experimental system.

## REFERENCES

al-Shamkhani A, Birkeland ML, Puklavec M, Brown MH, James W, Barclay AN. 1996. Eur J Immunol. Aug;26(8):1695-1699. Higgins LM, McDonald SA, Whittle N, Crockett N, Shields JG, MacDonald TT. 1999. J Immunol. Jan 1;162(1):486-493. Weinberg AD, Wegmann KW, Funatake C, Whitham RH. 1999. J Immunol. 162:1818-1826.

Tonbo Biosciences tests all antibodies by flow cytometry. Citations are provided as a resource for additional applications that have not been validated by Tonbo Biosciences. Please choose the appropriate format for each application and consult Materials and Methods sections for additional details about the use of any product in these publications.

For Research Use Only.

Not for use in diagnostic or therapeutic procedures. Not for resale. Not for distribution without written consent. Tonbo Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Tonbo Biosciences, Tonbo Biosciences Logo and all other trademarks are the property of Tonbo Biotechnologies Corporation. © 2013 Tonbo Biosciences.