

TECHNICAL DATA SHEET

Purified Anti-Mouse CD4 (GK1.5)

Catalog Number: 70-0041

PRODUCT INFORMATION

Contents: Purified Anti-Mouse CD4 (GK1.5)

Isotype: Rat IgG2b, kappa

Concentration: 0.5 mg/mL

Clone: GK1.5

Reactivity: Mouse

Formulation: 10 mM NaH₂PO₄, 150 mM NaCl, 0.09% Na₃N, pH7.2

DESCRIPTION

The GK1.5 antibody reacts with mouse CD4, a 55 kDa protein which acts as a co-receptor for the T cell receptor (TCR) in its interaction with MHC Class II molecules on antigen-presenting cells. The extracellular domain of CD4 binds to the beta-2 domain of MHC Class II, while its cytoplasmic tail provides a binding site for the tyrosine kinase lck, facilitating the signaling cascade that initiates T cell activation. CD4 is typically expressed on thymocytes, certain mature T cell populations such as Th17 and T regulatory (Treg) cells, as well as on dendritic cells. The GK1.5 antibody is widely used as a phenotypic marker for CD4 expression. If used together, the GK1.5 antibody and an alternative antibody, Anti-Mouse CD4 clone RM4-5, will compete for binding, i.e. RM4-5 is able to block GK1.5 binding to cells. In contrast, the Anti-Mouse CD4 clone RM4-4 does not block binding of the GK1.5 antibody to cells (Arora S et al. 2006. *Infect. Immun.* 74: 4339-4348). The GK1.5 antibody is also reported to be cross-reactive with Syrian hamster CD4.

PREPARATION & STORAGE

This monoclonal antibody preparation was purified from tissue culture supernatant via affinity chromatography. For In Vivo Ready™ (IVR) products, each preparation is also evaluated for endotoxin levels using the LAL assay. It is recommended to store the product undiluted at 4°C. Do not freeze.

APPLICATION NOTES

This purified format is guaranteed to be >90% pure as determined by SDS-PAGE analysis. Citations are provided as a convenience to you - please consult Materials and Methods sections for additional details about the use of any product in these publications.

REFERENCES

Taniguchi RT, DeVoss JJ, Moon JJ, Sidney J, Sette A, Jenkins MK, and Anderson MS. 2012. *Proc. Natl. Acad. Sci.* 109: 7847-7852. (in vivo depletion)Wolkers MC, Gerlach, C, Arens R, Janssen EM, Fitzgerald P, Schumacher TN, Medema JP, Green DR, and Schoenberger SP. 2012. *Blood.* 119:798-804. (in vivo depletion)Lee L-F, Logronio K, Tu GH, Zhai W, Ni I, Mei L, Dilley J, Yu J, et al. 2012. *Proc. Natl. Acad. Sci.* 10:1073. (flow cytometry).Thornton EE, Looney MR, Bose O, Sen D, Sheppard D, Locksley R, Huang X, and Krummel MF. 2012. *J. Exp. Med.* 10:1084. (immunohistochemistry – OCT embedded frozen tissue)Stephen TL, Wilson BS, and Laufer TM. 2012. *Proc. Natl. Acad. Sci.* 109: 7415-7420. (immunoprecipitation)Hammerbeck CD and Hooper JW. 2011. *J. Virol.* 85(19) 9929-9944. (flow cytometry – Syrian hamster).Diamond MS, Kinder M, Matsushita H, Mashayekhi M, Dunn GP, Archambault JM, Lee H, Arthur CD, White JM, Kalinke U, Murphy KM, and Schreiber RD. 2011. *J. Exp. Med.* 208: 1989-2003. (in vitro blocking)Kao H, Lin J, Littman DR, Shaw AS, and Allen PM. 2008. *J. Immunol.* 181: 8248-8257. (immunoprecipitation)Felix NJ, Donermeyer DL, Horvath S, Walters JJ, Gross M, suri A, and Allen PM. 2007. *Nat. Immunol.* 8:388-397. (in vitro blocking)

NOTE: Please choose the appropriate format for each application. Citations are provided as a convenience to you; please consult Materials and Methods sections for additional details about the use of any product in these publications.

For Research Use Only.

Not for use in diagnostic or therapeutic procedures. Not for resale. Not for distribution without written consent. Tonbo Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Tonbo Biosciences, Tonbo Biosciences Logo and all other trademarks are the property of Tonbo Biosciences Corporation. © 2013 Tonbo Biosciences.